

Hydrocarbon chemistry in inner regions of planet-forming disks

Jayatee Kanwar^{1,2,3}, Inga Kamp¹, P. Voitke^{2,3}, Ch. Rab⁴, W.F. Thi⁴

¹Kapteyn Astronomical Institute, University of Groningen, Groningen, The Netherlands

²Space Research Institute, Austrian Academy of Sciences, Graz, Austria

³TU Graz, Graz, Austria

⁴Max Planck Institute for Extraterrestrial Physics, Giessenbachstrasse, 85741 Garching, Germany




kanwar@astro.rug.nl

AIM: Study the carbon chemistry in the warm, dense inner regions of planet forming disks and its influence on the C₂H₂ spectrum observed by Spitzer and JWST.

Objectives:

- Study warm carbon chemistry in the inner 10au of T Tauri disks- nursery of planets
- Identify key destruction/formation pathways for C₂H₂ in the inner disk
- Study the impact on abundances and mid-infrared spectra.




JWST-MIRI spectrum of GW Lup[1] at R=3000


ProDiMo (Protoplanetary Disk Model)

The radiation thermo-chemical code ProDiMo [2] is used to model the warm chemistry in a standard T Tauri disk assuming the steady state and solar C/O.

The large DIANA chemical network [3] forms species up to C₄H⁺, the new extended chemical network has 92 species and can form hydrocarbons as large as C₈H₅⁺ and takes into account isotopomers. Ices of neutral species are included in both the networks. UMIST2012 and KIDA 2014 rate databases are used to calculate reaction rates.




The abundance of C₂H₂ using the extended chemical network. *Solid green:* emitting region of C₂H₂ at 13.7μm.




The distribution of the most abundant hydrocarbons in the inner disk using extended chemical network (excluding C and C⁺). The ices are represented by #.

Chemical Networks


The formation and destruction pathways for C₂H₂ in the region with T_{gas}= 270K and T_{dust}= 230K. The reaction rates are noted above the arrow as A(B) meaning Ax10^B.




The reactions for C₂H₂ found in KIDA 2014 missing from UMIST 2012:



Implications for Mid-Infrared Spectra



The flux emitted by C₂H₂ in the mid-infrared is modelled using FLiTs [4] at R=3000. The flux increases for the standard T Tauri disk by ~20% relative to the large DIANA network showing the effect of adding larger hydrocarbons.



The modelled C₂H₂ spectra using the extended chemical network with the UMIST 2012 and the KIDA 2014 database. The fluxes differ as each database leads to a different abundances. The T_{gas} is kept fixed in both models.

Future Work

Investigating chemistry for varying C/O ratios in disks and its affect on mid-IR spectra.

References

- [1] Grant, S. L., et al. 2023, A&A
- [2] Voitke, P., et al. 2018, A&A, 618, A57
- [3] Kamp, I., et al. 2017, A&A, 607, A41
- [4] Voitke, P., et al. 2018, A&A

